

Table 1. Characteristics of the all participants.

	Participants <i>n</i> (%)	Need for information <i>n</i> (%)
All	230 (100)	137 (59.6)
Gender		
Female	129 (56.1)	78 (60.5)
Male	100 (43.4)	58 (58.0)
Missing	1 (0.4)	
Age (years)	Median 64 (20-84)	
<50	24 (10.5)	12 (50.0)
50–70	153 (66.5)	98 (64.1)
>70	53 (23.0)	27 (50.9)
Cancer type*		
Prostate	66 (28.7)	45 (68.2)
Breast	56 (24.3)	40 (71.4)
Gastro-intestinal	46 (20.0)	28 (60.9)
Other	34 (14.8)	17 (50.0)
Not specified	36 (15.7)	14 (38.9)
Time since diagnosis		
≤2 year	51 (27.9)	36 (70.0)
3–5 year	72 (39.3)	48 (66.7)
6–10 year	39 (21.3)	21 (53.8)
>10 year	21 (11.5)	18 (56.3)
Stage of cancer		
Limited	151 (79.0)	97 (64.2)
Metastatic	38 (21.0)	24 (63.2)
Treatment*		
Surgery	157 (68.3)	98 (62.4)
External beam therapy	91 (39.6)	58 (63.7)
Chemotherapy	80 (34.8)	52 (65.0)
Hormonal therapy	54 (23.5)	36 (66.7)
Internal beam therapy	20 (8.7)	14 (70)

Self-reported need for information of the participants.

*Multiple answers possible.